

RESEARCH PAPER

Studying Toxigenic Fungi and Mycotoxins Detection from Bean (*Phaseolus vulgaris* L) in Duhok Province/Kurdistan of Iraq.

Lina F. Hussein ¹, Asia A. M. Saadullah ².

¹Department of Biology, College of Science, Duhok University- Duhok, Kurdistan Region, Iraq

²Department of Biology, College of Education, University of Duhok, Duhok, Kurdistan Region, Iraq

ABSTRACT:

The common bean (*Phaseolus vulgaris* L.) is one of the most widely farmed legumes, with farmers cultivating a variety of cultivar variants in Kurdistan of Iraq. Total of 50 bean samples were tested to establish their mycological contamination and their toxigenic potential under certain conditions, a variety of fungi may develop within bean grains; some of which have the capacity to synthesize mycotoxins. In current study toxigenic fungi were studied in beans (*Phaseolus vulgaris* L.) *Aspergillus* spp., *Fusarium* spp. and *Penicillium* spp. were the most frequently isolated genera, followed by *Phoma* sp., *Mucor* spp., *Alternaria* spp., *Curvularia* spp., *Rhizopus* spp., *Eurotium* sp. *Chaetomium* spp, Yeasts and *Drechslera* spp. Among 20 *Aspergillus* 11 strains produced mycotoxins: 30% produced aflatoxins (AFs) ranged between 81-260 ppb; 5% produced ochratoxin A (OTA 70-100 ppb) and 26.6% of *Fusarium* produced Trichothecene T-2/HT-2 (50- 94 ppb). The toxigenic species were *A. flavus*, *A. parasiticus*, *A. ochraceus*, *A. carbonarius* *A. niger* and *F. sporotrichioides*.

KEY WORDS: Toxigenic fungi, Phaseolus, Beans, Aflatoxins, Ochratoxin A, Trichothecenes.

DOI: <http://dx.doi.org/10.21271/ZJPAS.35.4.19>

ZJPAS (2023) , 35(4);192-199 .

1.INTRODUCTION :

Over the last three decades, food contamination by toxigenic molds has gotten a lot of attention. Filamentous and tiny fungus can infest most grains, including wheat, maize, bean, and rice. Some genera can create mycotoxins, which are harmful secondary metabolites that have an impact on food safety (Kuiper-Goodman, 1995).

Common bean (*Phaseolus vulgaris* L.) is a critical vegetable with tall wholesome esteem. Is one of the foremost historically developed crops within the world. It is of asocial, financial and dietary significance in numerous nations, utilized by at slightest 300 million individuals, developed significantly within the creating nations (Rangel *et al.*, 2005). A good source of protein is dried beans. carbohydrates and proteins, as well as vital vitamins, minerals, and fibers, and phenol compounds, which are rich in antioxidants.

Contamination of food and rural goods by various types of toxigenic molds could be a real and mostly overlooked problem. Despite decades of extensive research, form pollution continues to be a difficult problem (Gary Munkvold, 2017). FAO estimates that roughly 25% of the world's crops are contaminated by mold and impacted by mycotoxins with the loss that is expected amounting to billions of euros. Molds have been assigned a ranking as a source of harm in the capacity of grains, similar to scary crawly bugs. Inadequate collection of agricultural by-products and agricultural products such as beans, insufficient drying, handling, packaging, temperature, and excessive moisture during storage and shipping all contribute to the growth of fungi. and increase the risk of mycotoxin production, which are secondary metabolites given by toxigenic fungi in contaminated foods. Mycotoxins are released under fungal pressure, and their presence in a network indicates that the

* Corresponding Author:

Asia Abdulhamid Abdullah
Email: asia.saadullah@uod.ac

Article History

Received: 14/10/2022

Accepted: 05/01/2023

Published: 30/08 /2023

fungi were last seen lately (Sweeney and Dobson, 1999).

The problem of mycotoxins in food has resulted in increased concern about toxigenic fungal contamination, particularly from the genera *Aspergillus*, *Penicillium*, and *Fusarium*. These fungi have been discovered and identified from a variety of food substrates, including cereals, legumes and their by-products, and are widely distributed in the Duhok environment. Despite this, there are insufficient data on mycotoxins, fungal contamination, and toxigenic fungal potential in Duhok legumes, particularly beans and conditions of storage, this study was conducted to assess ochratoxin A (OTA) pollution and toxigenic contaminants, aflatoxins (AFs), and Trichothecenes 2 (T-2/HT-2) in beans using ELISA approach.

2. MATERIALS AND METHODS

2.1. MATERIALS

2.1.1 Samples collection and Mycoflora identification:

Fifty samples of beans (*Phaseolus vulgaris* L.), collected randomly at diverse time during one year (June 2020-June 2021) from general stores and road markets at Duhok Governorates. A 1,000 g minimum sample size was used. Tested Bean were partitioned into two bunches; the primary bunch was treated with sodium hypochlorite arrangement (1%) for two min (disinfected), whereas the second bunch was (untreated) on-disinfected. All tested beans were washed by sterilizing water for three minutes, then dried between two sterilized channel papers and plated on potato dextrose agar (PDA) and Dichloran Rose Bengal chloramphenicol (DRBC) agar. Ten beans/dish and three dishes were utilized as replication for each of the treatments. At 28°C, all plates were incubated for 5-7 days. All fungus development was exchanged and filtered single spore procedures onto PDA medium containing anti-microbial (chloramphenicol) to suppress bacterial growth (Klich, 2002).

Creating molds were refined on MEA medium (5-7 days incubation), then diagnosed at college of science, mycology research laboratory based on morphological and microscopic characteristics utilizing the accessible of writing concurring to Raper and Fennell (1965), Domsch et al. 1980, Klich, 2002 and Pitt 2009 for *Aspergillus*, Booth (1977) and Nelson et al. (1983) for *Fusarium*, and Singh et al. (1991) for

Aspergillus, *Penicillium* and *Fusarium*, see Maren and Johan (1988). On medium, the fungi's cultural features in terms of growth, colony character, texture, and sporulation were noted.

After seven days of incubation, measurements are taken. Lactophenol microscopic mounts with or without cotton blue are available. Microscopical slides were studied with oil immersion to identify the ornamentations of the stipe and conidia. Mycotoxins were extracted from fungal cultures, and the production of aflatoxins (AF), Ochratoxin A, and T2 by arbitrarily selected confines of *Aspergillus* section Flavi and section circumdati, as well as fusarium species, were screened using the Bragulat et al. (2001) used a centrally inoculating yeast extract sucrose (YES) plate method followed by a 7-day incubation at 25° C. At a halfway point between the edge and the middle of the forming colonies, agar plugs (0.5 cm in diameter) were evacuated from the colony's borders. In a tiny vial, the three plugs were combined with 1 ml methanol, forcefully shaken, and kept at room temperature for 1 hour before being blended again and the extricates separated. Via millipore channel (0.22um) width (Millex GP Channel Unit Carrigtwohill Co. Ireland). The following formula was used to calculate the isolation frequency of detected species from samples.

Isolation frequency FR % =

The number of samples on which a fungus has been found

Total number of samples

$\times 100$

2.1.2 Mycotoxins analysis

For analyses of mycotoxins the chemical connected immunosorbent test (ELISA) was used to examine the quantitative levels of AFs, OTA, and T-2/HT-2. The mycotoxins test was carried out in accordance with the information provided by the company (Veratox Aflatoxins, OTA, T2 quantitative Test, Neogen Corporation, USA). The number of mycotoxins produced by confinement was calculated using the standard bend obtained by mycotoxins measurements and expressed in parts per billion (ppb).

2.2 Analytical statistics

The data were evaluated using a one-way analysis of variance (ANOVA) followed by multiple mean comparisons and expressed as mean standard error (mean S.E). The data from this study was transformed to arcsine and analyzed using the SAS program, with Duncan's

multiple range test (SAS Institute Inc., Gary) was used to compare means.

3. RESULTS AND DISCUSSION

3.1. Extraction Yield

In Duhok province, the common bean is a widely cultivated legume. Agricultural products' temperature and moisture during storage and harvest allow fungi to grow, resulting in mycotoxins contamination, as well as their by-products, including beans, may be exposed to a high occurrence of Common beans contain biological contaminant., posing a food safety challenge because they can have an impact on the bean at various stages of the manufacturing process.

Their presence can have a negative impact and harm customers' health by having a direct harmful effect or by causing nutritional deficiencies, as well as lowering crop yields, which has an economic impact. The research literature on prevalence of biological contaminants in the common bean (*Phaseolus vulgaris* L.) was examined in order to determine the main concerns to food safety posed by fungal contamination and the possibility for mycotoxins.

Fungus contamination was investigated in 50 samples of beans gathered in Duhok province. A total of seventeen species from 13 genera were isolated. Table 1 and Figure 1 show the frequency of their occurrence.

Table 1: Frequency % of fungal genera on DRBC+PDA medium

Number	Frequency % of fungal genera on DRBC and PDA medium	
	Fungal genera	Frequency %
1	<i>Aspergillus parasiticus</i>	20.33
2	<i>A. flavus</i>	18.83
3	<i>A. niger</i>	18.5
4	<i>Fusarium sporotrichioides</i>	15.06
5	<i>A. carbonarius</i>	13.83
6	<i>A. fumigatus</i>	11
7	<i>Penicillium citrinum</i>	7.67
8	<i>Mucor circinelloids</i>	7
9	<i>A. foetidus</i>	6.17
10	<i>Rhizopus stolonifer</i>	6
11	<i>Alternaria alternata</i>	4.5
12	<i>A. nomius</i>	3.67
13	<i>A. ochraceus</i>	3.17
14	<i>Curvularia</i> sp.	3.17
15	<i>Phoma</i> sp.	2.83
16	<i>Eurotium amstelodami</i>	1.67
17	<i>Chaetomium</i> sp.	1.33
18	<i>P. expansum</i>	1.17
19	Yeasts nonidentified	1.06
20	<i>P. glabrum</i>	1
21	<i>Fusarium oxysporum</i>	1
22	<i>Drechslera</i> sp.	1




Figure 1: Frequency % of fungal genera on DRBC and PDA medium

Most of the recovered fungi were previously reported from beans in many parts of the world. Three species were isolated with high frequency namely: *Aspergillus parasiticus* (20.33), *A. flavus* (18.83) and *A. niger* (18.5), followed by *A. carbonarius* and *A. ochraceus* and *A. fumigatus* (13.83, 11,

3.17 %). Eight species of *Aspergillus* were found, with the most diversity of all of the retrieved taxa. When *A. parasiticus* and *A. flavus* grow on incorrectly stored grains, aflatoxins, one of the most dangerous carcinogens, are produced. *A. nomius*, in addition to black Aspergilli and Aspergilli in section Flavi, was found in this investigation on beans. *A. nomius* has been found in figs in California and has been reported to be aflatoxigenic (Bayman et al., 2002). Acute aflatoxicosis or, in severe cases, liver failure can result from ingesting greater quantities of aflatoxins (Fung et al., 2020).



Figure 2.a: Growth of *A. flavus* and *A. parasiticus* on Phaseolus bean on PDA medium.




Figure 2.b: Growth of *A. flavus* and *A. parasiticus* Phaseolus bean on DRBC medium.

Three kinds of black Aspergilli were found (*A. foetidus*, *A. carbonarius*, and *A. niger*). In the Kurdistan region, several species were detected in soil and other agricultural items (Abdullah, and Abdullah, 2009; Abdullah and Muhamed, 2011; Saadullah and Abdullah, 2012a, Abdullah and Saadullah, 2013).

Isolates from *A. carbonarius*, *A. ochraceus*, and to a lesser extent *A. niger* demonstrated production of ochratoxin A potential and were isolated often from dried vines naturally polluted with ochratoxin A in the Kurdistan region (Saadullah and Abdullah, 2012a). Because of their ability to create mycotoxins, *A. flavus*, *A. parasiticus*, *A. carbonarius*, and to a lesser extent, several species in the genus *Fusarium*, are the most important species contaminating beans (Moretti et al., 1998; and Heperkan et al., 2012a). *A. flavus* and *A. niger* were also the most frequent species on dried beans, according to Pitt and Hocking (2009).



Figure 3.a: Growth of *A. niger* on Phaseolus bean on PDA medium.




Figure 3.b: Growth of *A. carbonarius* and *A. niger* on Phaseolus bean on MEA medium.

Aspergillus spp. and *Penicillium* spp. are the most important storage fungus, while *Penicillium* spp. was represented by three species in the number of species isolated from beans. *P. citrinum* was the most frequent species (2.83%), followed by *P. expansum* and *P. glabrum* (1.17 and 1 %) respectively. Same results found by the Exudate Akwa et al. (2020) they isolated from beans in Kenya. There *Penicillium*, resistant species capable of growing at low water activity and tolerating high temperature (Smith and Onion, 1994).

These findings are comparable to those of Ruiz et al. (1996) in Argentina, who found that *Aspergillus* spp. and *Penicillium* spp. had the highest occurrence, with 40 and 15 percent, respectively, for freshly harvested beans. Scussel et al. (1998) looked at bean samples from various Brazilian states and found that these two genera were common (29.7 and 19.3 percent, respectively).

Fusarium sporotrichioides isolated in high frequency in the current study (15.06%) *Fusarium* is a phytopathogenic fungus that is one of the most economically important genera. This observation is concerning because *Fusarium* is a common food contamination that can lead to the development of fumonisins. *Fusarium langsethiae*, *F. Sporotrichioides*, and *F. Poae* are the most common *Fusarium* species. T-2 is rapidly converted in vivo to HT-2, which has comparable deleterious effects to T-2 with minor changes in strength (Schuhmacher-Wolz, U et al., 2016).



Figure 4.a: *Fusarium* on Phaseolus bean on PDA medium.



Figure 4.b: Growth of *A. ochraceus* on Phaseolus bean on MEA medium.

In addition, the high incidence of *Phoma* (7%) The presence of *Phoma* species in the beans analyzed poses a risk to consumers since some *Phoma* species can produce mycotoxins called cytochalasins (Griften, 1994).

The teleomorphic ascomycetes, namely *Eurotium amstelodami* were detected with percentage frequencies 1.67 %. Furthermore, when *E. amstelodami* isolates from Turkish dried figs were cultured on potato dextrose broth (PDB) medium, they produced Ochratoxin A and Aflatoxin B1 (Senyuva et al., 2008). *Eurotium* species can thrive in areas with limited water activity.

As a result, they're common in dishes with high sugar content (Pitt and Hocking, 1997). Other relatively frequent contaminants were, *Mucor*, *Rhizopus*, *Alternaria*, *Curvularia*, *Chaetomium*, yeasts and *Drechslera* Similar result was found by Romero et al. (2005).

Table 2 shows that from the total of *Aspergillus* strains isolated (20) from beans 30% were aflatoxigenic and 25% were ochratoxigenic. It revealed that the aflatoxins are more than

ochratoxins, while from the total *Fusarium* sp. 26.6% produced T2 this is in line with the study

done by (Costa et al., 1998 ,2000, 2002; Cruz et al., 1990).

Table 2: screening selected genera for aflatoxigenic, ochratoxigenic and T-2/HT-2 production abilities

Genera	N	Aflatoxins		Ochratoxins		T2/HT-2	
		N	%	N	%	N	%
<i>Aspergillus</i> sp.	20	6	30	5	25		
<i>Penicillium</i> sp.	10						
<i>Fusarium</i> sp.	8					3	26.6
Others	30						

Table 3: Quantitative production of Aflatoxins and Ochratoxin A and T2 by *Aspergillus* and

***Fusarium* species in vitro by ELISA technique.**

Fungal sp.	AFs ppb	OTA ppb	T2 ppb
<i>A. flavus</i> 1	260.0		
<i>A. parasiticus</i> 1	200.0		
<i>A. parasiticus</i> 2	145.0		
<i>A. parasiticus</i> 3	100.0		
<i>A. flavus</i> 2	81.0		
<i>A. ochraceus</i>		73	
<i>A. carbonarius</i>		100	
<i>A. niger</i>		70	
<i>Fusarium sporotrichioides</i> 1			50
<i>Fusarium oxysporum</i> 2			94
<i>Fusarium oxysporum</i> 3			58

Table 3 showed the results of screening four isolates for each of *A. flavus*, and *A. parasiticus* *A. ochraceous*, *A. carbonarius* and *A.*

niger, *Fusarium* isolated from dried beans for their aflatoxigenic, ochratoxigenic and T-2/HT-2 production abilities in culture media as detected by ELISA technique. Two strains of *A. flavus*

showed positive abilities as aflatoxin producers, whereas Three *A. parasiticus* were positive, and *A. ochraceus*, *A. carbonarius* and *A. niger* to produce OTA. The initial strategy to controlling the incidence of aflatoxins and ochratoxins in beans was to control contamination with aflatoxins and OTA in the field, which is extremely difficult to do because it is predominantly driven by meteorological variables like relative humidity

and temperature (Applebaum et al., 1982).

The highest levels of these three mycotoxins, on the other hand, have been linked to the post-harvest growth of *Aspergillus* and *Fusarium* molds on improperly preserved foods (Jay, 1992).

The aflatoxin, OTA, and T-2/HT-2 potential of the isolates studied varied significantly. This is consistent with several other studies that show that the ability to produce aflatoxin varies among *Aspergillus* section flavi isolates (Senyva et al., 2008, Embaby et al., 2012). According to various studies, not all strains of *A. flavus* may produce aflatoxin, and the ratio of non-aflatoxigenic to aflatoxin-producing isolates varies depending on the isolates' source and location (Schroeder and Bolla 1973; Abdel-Malik et al., 1993; Abdullah and Al-Mousawi, 2009; Abdullah et al., 2009).

4. CONCLUSIONS

In conclusion, based on our results that the contamination of beans by aflatoxigenic strains of *A. flavus* and *A. parasiticus* and by ochratoxigenic strains of *A. carbonarius* as well as by trichothecenes 2 of *Fusarium sporotrichodes* and *F. oxysporum* T-2/HT-2 represent a special hazard to consumers health. Therefore, such contaminated materials should be monitored before use.

References

- ABDEL-MALEK, A.Y., EL-MARAGHY, S.S AND HASSAN, H.A.A.1993, Mycotoxin-producing potential of some *Aspergillus*, *Penicillium* and *Fusarium* isolates found on corn grains and sunflower seeds in Egypt. *Journal Islamic Academic.Science*.6:189-192.
- ABDULLAH, S. K. AND MOHAMMED, A. A. 2011, Prevalence of Black *Aspergilli* in soil at vineyard in Duhok, Kurdistan Region of Iraq, *Proc. 4th Int. Sci. Con. Salaheddin Univ. Erbil*. October 18-20, 2011, vol.3 pp966-969.
- ABDULLAH, S.K. AND AL-MOUSAWI, K.A. 2009, Incidence of *Aspergillus* species in seeds of corn and sunflower cultivars grown in Iraq and aflatoxin producing potential of *Aspergillus* section Flavi. *Proc.1st Scient. Conf.Biol.Sci.Mosul University*, 22-23 April 2009, P299-307.
- ABDULLAH, W.R. AND ABDULLAH, S.K. 2009, Taxonomic study on black *Aspergilli* from soil in Kurdistan region of Iraq. *Journal of Duhok University*. 12 (special issue), 288-295.
- ABDULLAH, S.CK AND ASADULLAH. 2013, Soil mycobiota at grapevine plantations in Duhok, North Iraq. *Mesopotamia J.Agric*.41 (Suppl.1):402-412.
- APPLEBAUM RS, BRACKETT RE, WISEMAN DW, MARTH EH. 1982, Aflatoxin: toxicity in dairy cattle and occurrence in milk and dairy products-a review. *Journal of Food Protect* ,45:752-777.
- BAYMAN, P., BAKER, J.L., DOSTER, M.A., MICHAILIDES, T.J AND MAHONEY, N.E. 2002, Ochratoxin production by *Aspergillus ochraceus* group and *Aspergillus alliaceous*. *Applied Environmental Microbiology* 68: 2326-2329. *Biol. Sci. University of Duhok* 8-10 May, 2012 pp. 260-268.
- BOOTH, C. 1977, The genus *Fusarium*. First published. In *commonwealth Mycological Institute, Kew, Surrey, England* pp. 235.
- BRAGULAT, M.R., ABARCA, M.L., CABANES, F.J., (2001). An easy screening method for fungi producing ochratoxin A in pure culture. *International Journal of Food Microbiology*. 139-144.
- COSTA Costa, L.L.F.; Scussel, V.M. Toxigenic fungi in beans (*Phaseolus vulgaris* L.). 1998, *Revue de Medecine Veterinaire*, p.518, fresh and dry fruit quality contaminated by some mold fungi *Journal of Applied Science. Res.*, 8:3083-3091 from Brazil. *Revue de Medecine Veterinaire*, 1998. p.531.
- COSTA; LÉA LUZIA FREITAS AND SCUSSEL VILDES MARIA 2002, Toxigenic fungi in Beans (*Phaseolus vulgaris* L.) classes black and colour cultivated in the state of Santa catarina Brazil. *Brazilian. Brazilian Journal of Microbiology* (2002) 33:138-144 ISSN 1517-8382
- CRUZ, L.C.H.; SANTUÁRIO, J.M.; ROSA, C.A.R.; CAMPOS,S.G. 1990, Caracterização da microbiota toxigênica da superfície do feijão VI Encontro Nacional de Micotoxinas. São Paulo. p.57.
- DOMSCH, K. H., GAMS, W. AND ANDERSON, T. H. 1980, *Compendium of soil fungi*. Academic Press, London.
- EMBABY, E.M.; HAGGAG, L.F.; ABDEL-GALIL, M. 2012, Decay of some fresh and dry fruit quality contaminated by some mold fungi. *Journal Applied of Science. Res.* 8, 3083-3091.
- EXODUS AKWA.JOHN MAINGI, JONAH KIPRONO BIRGEN. 2020, Assessing the

- incidence of common bean (*Phaseolus vulgaris* L.) infection by fungi during harvest and storage in Cameroon, DOI: 10.5943/sif/5/1/24.
- FUNG F., Clark R. and Williams. 2002, *Stachybotrys*, a Mycotoxin-producing Fungus of Increasing Toxicogenic Importance, *Journal Toxicology, Clinical Toxicology*,79-86.
- GARY MUNKVOLD 2017, "In book: "Mycotoxigenic Fungi." *Fusarium* species and their associated mycotoxins. *Methods in molecular biology* (Clifton, N.J.) 1542:51-10 DOI: 10.1007/978-1-4939-6707-0_4.HATAMI, T., EMAMI, S. A., MIRAGHAEI, S. S. & MOJARRAB, M. 2014. Total phenolic contents and antioxidant activities of different extracts and fractions from the aerial parts of *Artemisia biennis* Willd. *Iranian journal of pharmaceutical research: IJPR*, 13, 551.
- GRIFFIN, D.H. 1994, *Fungal Physiology*, 2nd ed. New York: Wiley-Liss, 458p.
- HEPERKAN, D., KARBANCIOGLA-GULER, F. AND OKTAY, H. I. 2012a, Mycoflora and natural occurrence of aflatoxin, cyclopiazonic acid, fumonisin and ochratoxin A in dried figs. *Food Additives and Contaminates*. 29:277- 286.
- SAMSON, R.A. HOEKSTRA J. A.M AND. FRISVAD., J.C. 2004, *Introduction to Food and Airborne Fungi*. 7th Edition. Centraal bureauvoor Schimmel cultures, Utrecht.
- JAY JM. 1992, *Modern food microbiology*, 4th edn. Champman and Hall, New York, pp 1– 70.
- KLICH, M.A. 2002, Identification of common *Aspergillus* species. *CentralbureauVoor*.
- KUIPER-GOODMAN, (1995). *Mycotoxins: risk assessment and legislation*. *Toxicol Lett* 82(83):853–859.
- MAREN, A.K. AND I.P. JOHAN 1988, *A Laboratory guide to the common Aspergillus spp. and their teleomorph*. *Commonwealth Scientific and Industrial*, pp. 116.
- MICHAEL J. SWEENEY, ALAN D.W. DOBSON. 1999, *Molecular biology of mycotoxin biosynthesis*. *FEMS Microbiology Letters* 175, 149-163
- MORETTI A, MULE G, PERRONE G AND LOGRIECO A, 1998, Genetic and toxigenic characterization of *Fusarium moniliforme* and *Fusarium proliferatum* from various plants. In *Sixth International Mycological Congress* (pp. 166). Jerusalem, Israel.
- NELSON TOUSSOUN AND MARASAS 1983, *Fusarium spp. An Illustrated Manual for Identification*. Published by the Pennsylvania state university.
- PITT, J. I. AND A. D. HOCKING 1997, *Fungi and Food Spoilage* 2nd ed. Blackie Academic & Professional. London, U.K. pp.511.
- PITT, J. I. AND HOCKING, A. D. 2009, *Fungi and food spoilage* 3rd edition. Springer, New York, N. Y. USA, 540pp. potential of *Aspergillus* strains isolated from vineyard soil, fresh grape berries potential of some *Aspergillus*, *penicillium* and *Fusarium* isolates found on corn grains and sunflower seeds in Egypt. *Journal of Islamic Academic.Sci*.6:189-192.
- RANGEL AF, MOBIN M, RAO IM, ET AL. 2005, Proton Toxicity Interferes With The Screening Of common bean (*Phaseolus vulgaris* L.) genotypes for aluminum resistance in nutrient solution. *Journal of Plant Nutrition and Soil Science* ,168: 607-616.
- MAZHAR, A. 2014. RAPER, K.B.; FENNELL, D.I. 1965, *The Genus Aspergillus*. Baltimore. Williams & Wilkins Company, 686p.
- ROMERO, S.M., COMERIA, R.M., LARUMBA, G., RITIENI A., VAAMNDA, G. AND PINTO, V.F. 2005, Toxigenic fungi isolated from dried vine fruits in Argentina *Int. Journal of Food Microbiology.*, 104: 43-49.
- RUIZ, J.A.; BENTABOL, A.; GALLEGO, C.; ANGULO, R.; JODRAL, M. 1996. Mycoflora and aflatoxin-producing strains of *Aspergillus flavus* in greenhouse-cultivated green beans (*Phaseolus vulgaris* L.). *Journal of Food Protection*, 59(4): 433-435.
- SAADULLAH, A. A. M AND ABDULLAH, S. K. 2012a, New records of ochratoxin producing species of *Aspergillus* contaminated raisins from Duhok, Kurdistan region, Iraq. *Journal of University of Duhok* .15 (special issue): 377-389.
- SAS 1999, *SAS/STAT User's Guide*, Version 8.2, 1st printing. Vol. 2. SAS Institute Inc, SAS Campus Drive, Cary, North Carolina.Schroeder, H. W. and Bolla, R. A. 1973, *Food Mycology*.
- SCHUHMACHER-WOLZ, U.; HEINE, K.; SCHNEIDER, K.REPORT 2016, on Toxicity Data on Trichothecene Mycotoxins HT-2 andT-2 Toxins. CT/EFSA/CONTAM/2010/03. Available online: http://www.efsa.europa.eu/en/scdocs/scdoc/65_e.htm (accessed on 9 April 2016).
- SENYUVA, H.Z, GILBERT, J., SANESON, R.A., OZCAN, S., OZTURKOGLU, S. AND ONAL, D. 2008, Occurrence of fungi and their mycotoxins in individual Turkish dried figs. *Journal of World Mycotoxin* .1:79-86.
- SINGH, K.; C. JENZ, U. THRON AND S.B. MATHUR 1991, *An Illustrated of some Seed-born Aspergilli, Fusaria, Penicillia and their mycotoxins*. First edition, Danish Government Institute of Seed Pathology for Developing Countries. Ryvangs Alle 78. DK-2900 Hellerup, Denmark and Department of Biotechnology.
- SMITH, D. AND ONIONS, A.H.S. (1994). *The Preservation and Maintenanc of Living Fungi*, 2nd. *International Mycological Institute*. pp. 17–20. Wallingford, UK: CAB International.
- SWEENEY MICHAEL J., DOBSON ALAN D.W. 1999, *Molecular biology of mycotoxin biosynthesis*. *FEMS Microbiology Letters* 175 (1999) 149 -163.